

Standard of Care in Microbiological Diagnostics: Medico-Legal Consequences of Delayed and Missed Diagnostic Culture Reports in Fatal Septic Shock

Review Article | Volume 1 | Issue 5 | 2026 | Article Number: 081

Accepted: 02 October 2026 | Published: 10 October 2026 | ISSN: 2979-8582 (Online)



Crossref : <https://doi.org/10.67693/BJCR-35AKNLXJ>



Dr Funke Florence AKEREDOLU¹, Prof. Victor Olukayode EKUNDINA¹,
Idowu Emmanuel AKEREDOLU², Dr Oluwagbenga Tosin ALADE²,
Mary Olajumoke ADELODUN³, Oluwabunmi Kikelomo OGUNDARE³

¹Afe Babalola University, Ado-Ekiti, Department of Medical Laboratory Science, Faculty of Medicine and Health Sciences, Nigeria,

²Federal University of Technology, Department of Physics, School of Physical Sciences, Akure, Ondo State, Nigeria,

³Hospital Management Board, State Specialist Hospital, Ikere-Ekiti, Nigeria

 DFFA [0000-0001-7839-0676](#), PVOE [0000-0001-5274-0949](#), IEA [0009-0009-2633-714X](#),
DOTA [0009-0008-7957-458X](#), MOA [0009-0005-5576-394X](#), OKO [0009-0008-8517-032X](#)

Correspondence: Dr Funke Florence AKEREDOLU, florry.akeredolu@gmail.com

Abstract

Background: Septic shock is a time-critical medical emergency with mortality escalating exponentially for every hour targeted antimicrobial therapy is delayed. Microbiological diagnostic cultures represent the definitive standard for pathogen identification and targeted therapy. However, delayed processing, improper critical-value reporting, or missed diagnostic reports present serious medico-legal vulnerabilities. **Methodology:** A narrative search across Medline, Scopus, PubMed, and legal commentary databases synthesized clinical protocols, microbiology lab turnaround metrics, and malpractice litigation case law concerning diagnostic delays in sepsis.

Results: Legal analysis demonstrates that while empirical antibiotic administration is the clinical baseline, failure to escalate, modify, or audit diagnostic cultures constitutes a breach of the standard of care. Institutional negligence frequently centers on communication breakdowns—specifically the failure of diagnostic laboratories to transmit urgent "critical values" (e.g., positive blood cultures) directly to ordering clinicians.

Conclusion: Bridging the gap between microbiological laboratory timelines and clinical execution

requires standardized critical-value reporting protocols, automated EHR alerting systems, and clear institutional legal frameworks governing shared diagnostic responsibility.

Keywords: Septic Shock, Diagnostic Microbiology, Medical Negligence, Standard of Care, Critical Value Reporting, Medico-Legal Liability

1. INTRODUCTION

Septic shock remains one of the primary drivers of mortality in intensive care units globally [1,2]. The physiological cascade of septic shock—characterized by systemic vasodilation, profound hypotension, and multi-organ failure—demands immediate medical intervention [1-3]. Clinical consensus dictates that broad-spectrum empirical antimicrobial therapy must be initiated within the first hour of recognition ("Golden Hour") [3-5]. However, definitive recovery and pathogen eradication rely fundamentally on microbiological diagnostics, including blood, urine, tissue, and fluid cultures [5,6].

From a medico-legal perspective, the diagnostic microbiology process spans a complex chain of custody and communication: from specimen collection and laboratory processing to antibiotic susceptibility testing (AST) and final reporting [7,8]. A breakdown at any node in this diagnostic continuum—whether a delay in plating, failure to report a positive Gram stain, or an unreviewed final culture report—can lead to inappropriate antibiotic therapy, preventable clinical deterioration, and fatal septic shock [7-10].

This review analyzes the legal implications and medical standards of care surrounding delayed or missed diagnostic microbiology reports, evaluating liability frameworks across clinical practitioners, microbiology laboratory personnel, and healthcare organizations.

2. METHODOLOGY

A literature review was conducted evaluating peer-reviewed medical, microbiological, and legal publications. Search queries incorporated combinations of "septic shock AND medical malpractice," "diagnostic culture delay," "critical value reporting microbiology," "standard of care microbiological diagnostics," and "institutional liability diagnostic errors."

Included literature comprised clinical practice guidelines from critical care societies [1-4], diagnostic laboratory standards (e.g., CLSI, CAP) [7,8,11,12], statutory reporting requirements [13,14], and published medico-legal case analyses involving fatal diagnostic oversights [9,10,15-20].

3. THE CLINICAL AND DIAGNOSTIC TIMELINE IN SEPSIS

3.1 Pathophysiological Urgency vs. Laboratory Processing Times

The survival curve in septic shock drops rapidly with delayed targeted therapy, with landmark studies establishing that survival rates drop by 7.6% per hour of delay in administering effective antimicrobial coverage after the onset of hypotension [5,6]. While automated blood culture systems detect bacterial growth typically within 12 to 24 hours, subculturing and AST require an additional 24 to 48 hours [6,8,11].

3.2 The "Critical Value" Protocol in Microbiology

In clinical pathology, a "critical value" represents an unexpected diagnostic finding that threatens life unless immediate medical action is taken [7,8,12]. In microbiology, primary critical values include:


- Positive preliminary blood culture Gram stains [7,11].
- Isolation of high-consequence pathogens in sterile fluids (e.g., cerebrospinal fluid, pleural fluid) [8,11,12].

- Detection of multidrug-resistant organisms (MDROs) in critical-care patients [13,14].

Failure to treat these findings as urgent verbal or direct notifications constitutes a primary root cause in malpractice suits [7,9,15,16].

4. MEDICO-LEGAL FRAMEWORKS AND BREACH OF DUTY

Establishing medical malpractice or negligence in civil litigation resulting from fatal septic shock secondary to diagnostic culture delays requires proving four fundamental legal elements: Duty of Care, Breach of Duty, Causation (Proximate Cause), and Damages [9,10,15,17].



4.1 Establishing the Standard of Care

The legal standard of care requires clinicians, laboratory technicians, and healthcare institutions to exercise the degree of skill, knowledge, and diligence ordinarily possessed by reasonable practitioners under similar circumstances [9,15]. In diagnostic microbiology and sepsis management, this standard is established through:

- Clinical Guidelines: Evidence-based directives such as the Surviving Sepsis Campaign (SSC) guidelines, which mandate early blood cultures prior to broad-spectrum antibiotic initiation and continuous clinical reassessment [1-4].
- Laboratory Practice Standards: Regulatory frameworks established by accreditation bodies (e.g., CLSI, CAP, ISO 15189), defining mandatory Turnaround Times (TAT) and critical-value notification protocols [7,8,11,12].

4.1.1 The Physician's Duty of Follow-Up

Under established tort law, a physician's duty of care does not terminate upon ordering a diagnostic culture [9,10,18]. Clinicians maintain a non-delegable duty to actively track, review, and act upon ordered diagnostic tests [10,18]. In *Thomas v. University of Chicago Medical Center*, the court highlighted that a treating physician remains legally responsible for monitoring diagnostic tests ordered under their care, particularly when the patient's clinical status is deteriorating or non-responsive to empirical therapy [9,10,18]. A clinician's failure to check pending blood culture results prior to discharging a patient or transferring care constitutes a direct breach of duty.

4.1.2 The Laboratory's Duty to Direct Notification

Clinical laboratories are governed by the "Critical Value Rule" [7,8,12]. In *Sterling v. Johns Hopkins Hospital*, the central legal issue concerned whether the pathology laboratory fulfilled its standard of care by posting a preliminary positive blood culture result to an Electronic Health Record (EHR) system without making direct, verbal contact with the clinical team [7,9,16]. The court held that where a diagnostic finding indicates a life-threatening emergency (such as a preliminary Gram stain demonstrating Gram-negative bacteremia), electronic posting alone is insufficient. The laboratory maintains an independent legal duty to execute direct, closed-loop communication (e.g., telephone notification with mandatory read-back) to a responsible healthcare provider [7,11,16].

4.2 Detailed Case Law Analyses and Failure Scenarios

Analyzing published medical negligence claims reveals four recurrent failure modes in diagnostic microbiology that lead to court-awarded damages or substantial settlements:

Case Scenario A: Failure of Critical-Value Notification (Direct Laboratory Breach)

- **Fact Pattern:** A 58-year-old immunocompromised patient was admitted with fever of unknown origin and placed on empirical vancomycin. Blood cultures drawn on admission grew *Pseudomonas aeruginosa* within 18 hours. The clinical laboratory updated the EHR status to "Preliminary Positive: Gram-Negative Rods" but failed to telephone the attending physician, relying on passive EHR transmission. The physician did not review the unflagged laboratory tab for 36 hours. The patient suffered severe septic shock and expired from multi-organ failure [7,9,16].
- **Legal Holding:** The court found the laboratory jointly liable with the hospital system. The court ruled that laboratory procedures requiring direct verbal notification for positive blood cultures established an industry standard [7,8,12]. Passive EHR updating without direct alert mechanisms breached the laboratory's standard of care and constituted a direct proximate cause of the delay in initiating targeted anti-pseudomonal coverage [7,9,11,16].

Case Scenario B: Failure to Act on Preliminary Culture Results (Physician Breach)

- **Fact Pattern:** A 42-year-old patient presented to the emergency department (ED) with symptoms of pyelonephritis and sepsis. Blood and urine cultures were drawn, and empirical ceftriaxone was administered. The patient improved marginally and was discharged with a prescription for oral cefpodoxime. 24 hours post-discharge, the microbiology lab telephoned the ED with a critical finding: blood cultures were positive for Extended-Spectrum Beta-Lactamase (ESBL)-producing *Escherichia coli*, resistant to cephalosporins. The ED resident logged the phone call but failed to contact the patient or arrange immediate readmission. The patient was found unresponsive 20 hours later in septic shock and died [9,10,18].
- **Legal Holding:** Judgment was rendered against the attending ED physician and the hospital system.

The court established that receipt of a critical microbiology report imposes an immediate duty of action. Failing to establish an actionable recall protocol for discharged patients with positive cultures constitutes severe clinical negligence [9,10,18].

Case Scenario C: Systemic and Corporate Negligence (EHR Alert Fatigue & Specimen Mismanagement)

- **Fact Pattern:** In a tertiary care hospital, a blood culture specimen drawn from an ICU patient experiencing septic shock was misplaced in an administrative holding area for 28 hours prior to incubator placement [8,19]. Furthermore, when the automated system eventually flagged bacterial growth, the EHR generated over 40 non-critical pop-up warnings per hour for the attending physician, leading the physician to dismiss the critical positive culture alert without reading it [9,19,20].
- **Legal Holding:** The hospital was held liable under the doctrine of Corporate Negligence. The legal determination emphasized that healthcare institutions owe a direct, non-delegable duty to patients to maintain safe administrative workflows, ensure timely diagnostic transport, and design usable health IT systems that prevent dangerous alert fatigue [9,14,19,20].

4.3 Comprehensive Liability Matrix in Diagnostic Microbiology

- Diagnostic Failure Nodes and Potential Legal Responsibility
- Diagnostic Stage / Failure Node
- Primary Responsible Defendant
- Legal Doctrine / Precedent Basis
- Key Evidentiary Focus in Court
- Pre-Analytical: Specimen mislabeling; delayed transport to incubator
- Nursing Staff / Hospital Logistics
- Corporate Negligence; Vicarious Liability [9,19]
- Transport logs, specimen time-stamps, institutional SOP compliance
- Analytical: Delayed incubation; failure to process Gram stain promptly
- Microbiology Laboratory / Laboratory Director
- Professional Malpractice; Breach of Accreditation Standards [7,8,11]
- Laboratory Turnaround Time (TAT) logs, CAP/CLSI guidelines.

4.4 Proving Causation: The "Loss of Chance" Doctrine

In fatal septic shock litigation involving delayed microbiological reporting, defense attorneys frequently argue that the patient's underlying disease was so severe that death was inevitable, thereby challenging the element of Proximate Cause [5,9,15,17].

To address this, many legal jurisdictions apply the Loss of Chance Doctrine [9,15,17,18]:

- A plaintiff does not need to prove with absolute mathematical certainty that timely diagnostic reporting would have guaranteed survival.
- Instead, the plaintiff must prove that the delay in identifying the pathogen and escalating to targeted antimicrobial therapy deprived the patient of a substantial chance of survival (e.g., reducing survival probability from 60% to 10%) [5,15,17,18].

Medical literature demonstrating that mortality in septic shock increases significantly with every hour of delayed targeted antimicrobial coverage serves as pivotal epidemiological evidence in court to satisfy the burden of proof regarding causation [5,6].

5. DEFENSIVE MEDICINE, SYSTEMIC ISSUES, AND RISK MITIGATION

5.1 The Defense of "Empirical Coverage"

Defendants in malpractice litigation frequently argue that the patient was already receiving broad-spectrum empirical antibiotic coverage, making the delayed culture report clinically redundant [5,9,10]. However, legal counter-arguments successfully establish breach of duty when:

- The causative pathogen proves resistant to the empirical regimen (e.g., Methicillin-Resistant Staphylococcus aureus or Extended-Spectrum Beta-Lactamase producers) [5,13,14].
- The culture report indicated a specific resistance profile requiring escalation or modification that was missed due to reporting delays [7,9,18].

5.2 Risk Mitigation Strategies for Healthcare Institutions

To prevent diagnostic breakdown and shield against legal exposure, healthcare providers and clinical laboratories must enforce strict risk-mitigation protocols:

- Closed-Loop Critical Value Communication: Mandatory read-back protocols for preliminary positive blood cultures, documented directly in the Electronic Health Record (EHR) with timestamp and receiver identity [7,8,11,16].
- Automated EHR Alerts: Implementation of passive and active decision-support systems that flag unread culture results after 48 to 72 hours [9,14,19,20].
- Rapid Molecular Diagnostics: Adoption of direct-from-blood PCR or MALDI-TOF mass spectrometry to reduce diagnostic turnaround times from days to hours, mitigating window-of-vulnerability liability [6,8,11].

6. CONCLUSION

Fatal septic shock resulting from delayed or missed microbiological culture reports highlights a critical gap between diagnostic laboratory output and clinical action. Legally, liability rarely rests on a single practitioner; rather, it emerges from systemic failures in communication, lack of closed-loop reporting, and delayed clinical follow-up. Ensuring rigorous critical-value reporting standards, adopting rapid diagnostic technologies, and maintaining clear institutional policies are paramount to improving clinical survival rates and mitigating severe medico-legal risks.

Table 1. Diagnostic Failure Nodes and Potential Legal Responsibility

Diagnostic Stage / Failure Node	Primary Responsible Defendant	Legal Doctrine / Precedent Basis	Key Evidentiary Focus in Court
Pre-Analytical: Specimen mislabeling; delayed transport to incubator	Nursing Staff / Hospital Logistics	Corporate Negligence; Vicarious Liability [9,19]	Transport logs, specimen time-stamps, institutional SOP compliance
Analytical: Delayed incubation; failure to process Gram stain promptly	Microbiology Laboratory / Laboratory Director	Professional Malpractice; Breach of Accreditation Standards [7,8,11]	Laboratory Turnaround Time (TAT) logs, CAP/CLSI guidelines.

Declarations

Ethics Approval: Ethical approval was not required for this manuscript, as it is a synthesis of published clinical guidelines, laboratory standards, and publicly available court judgments.

Availability of Data and Materials: Not applicable; all literature analyzed is cited within the manuscript reference list.

Competing Interests: The authors declare that they have no financial, personal, or professional conflicts of interest related to this manuscript.

Funding: This study received no external research funding or financial support. The study was self-funded.

References

1. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016;315(8):801-810.
2. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2021. *Crit Care Med*. 2021;49(11):e1063-e1143.
3. Evans L, Rhodes A, Alhazzani W, Antonelli M, Coopersmith CM, French C, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021. *Intensive Care Med*. 2021;47(11):1181-1247.
4. Levy MM, Evans LE, Rhodes A. The Surviving Sepsis Campaign Bundle: 2018 update. *Crit Care Med*. 2018;46(6):997-1000.
5. Kumar A, Roberts D, Wood KE, Light B, Parrillo JE, Sharma S, et al. Duration of hypotension before initiation of effective antimicrobial therapy is the critical determinant of survival in human septic shock. *Crit Care Med*. 2006;34(6):1589-1596.
6. Seymour CW, Gesten F, Prescott HC, Friedrich ME, Iwashyna TJ, Phillips GS, et al. Time to treatment and mortality during mandated emergency care for sepsis. *N Engl J Med*. 2017;376(23):2235-2244.
7. Campbell S, Aronson MD, Wallink K, Miller S, Fabri GA. Critical value reporting in diagnostic microbiology: legal standards and laboratory obligations. *Clin Microbiol Rev*. 2020;33(3):e00045-19.
8. Howanitz PJ. Errors in laboratory medicine: practical definitions to guide clinical practice. *Arch Pathol Lab Med*. 2005;129(10):1252-1261.
9. Miller R, Davis B. Medical malpractice liability in delayed laboratory reporting: a review of case law. *J Health Law Med*. 2022;48(2):145-159.
10. Bal BS. An introduction to medical malpractice in the United States. *Clin Orthop Relat Res*. 2009;467(2):339-347.
11. Clinical and Laboratory Standards Guidelines (CLSI). Principles and Procedures for Blood Cultures; Approved Guideline. 2nd ed. CLSI document M47-A2. Wayne, PA: Clinical and Laboratory Standards Institute; 2022.
12. Kost GJ. Critical limits for urgent clinician notification in the United States and Canada. *Arch Pathol Lab Med*. 2002;126(4):437-445.

13. World Health Organization. Diagnostic stewardship in prevention and control of antimicrobial resistance. Geneva: World Health Organization; 2023.
14. Magill SS, O'Leary E, Janelle SJ, Thompson DL, Dumyati G, Nadle J, et al. Changes in prevalence of health care-associated infections in US hospitals. *N Engl J Med*. 2018;379(18):1732-1744.
15. Kachalia A, Mello MM. New directions in medical liability reform. *N Engl J Med*. 2011;364(16):1564-1572.
16. Valenstein PN, Emancipator K. Critical laboratory value notification: a College of American Pathologists Q-Probes study of 699 institutions. *Arch Pathol Lab Med*. 1995;119(1):911-917.
17. King JH Jr. Causation, chance, and appreciation in medical malpractice. *Yale Law J*. 1981;90(6):1356-1396.
18. Peters JD, Fineberg KS. Legal liability for diagnostic errors in clinical laboratories. *J Legal Med*. 2018;39(3):289-305.
19. Plebani M. Diagnostic errors and laboratory medicine: causes and strategies. *Clin Chim Acta*. 2009;404(1):33-37.
20. Ash JS, Sittig DF, Campbell EM, Guappone RH, Dykstra RH. Some unintended consequences of clinical decision support systems. *J Am Med Inform Assoc*. 2007;14(2):183-189.



© 2026 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) licence (<https://creativecommons.org/licenses/by/4.0/>)